

## VACCINATION MINIMUM INTERVALS

VACCINE	AGE	MINIMUM INTERVAL	NOTES
IPV (POLIO) #1	2 MONTHS	4 WEEKS	4 Polio given at any
IPV #2	4 MONTHS	4 WEEKS	age is the maximum
IPV #3	6 MONTHS	4 WEEKS	
IPV #4	4 TO 6 YEARS	-	
DTaP #1	2 MONTHS	4 WEEKS	For school, if 4 <sup>th</sup>
DTaP #2	4 MONTHS	4 WEEKS	DTaP given <u>≥</u> 4
DTaP #3	6 MONTHS	6 MONTHS	birthday, series
DTaP #4	12 TO 15 MONTHS	6 MONTHS	Complete (series
DTaP #5	4 TO 6 YEARS	-	complete is 5 doses).
Hib # 1/COMVAX	2 MOS.	4 WEEKS	Comvax cannot be
Hib #2/COMVAX	4 MOS.	4 WEEKS	administered to
(Hib #3) +!	(6 MOS.)	8 WEEKS	Infants <6 weeks of
Hib #4/COMVAX	12 TO 15 MOS.	-	age. *
MMR #1	12 – 15 MOS.	4 WEEKS	
MMR #2	4 – 6 YEARS	-	
HEP B #1 !	BIRTH / 2 MOS.	4 WEEKS	Infant cannot finish
HEP B #2	1 MOS. - 4 MOS.	8 WEEKS	series prior to age
HEP B #3 !!	6 MOS. - 18 MOS.	-	6 mos.
PREVNAR #1(PCV7)	2 MOS.	4 WEEKS	Refer to table
PREVNAR #2	4 MOS.	4 WEEKS	
PREVNAR #3	6 MOS.	8 WEEKS	
PREVNAR #4	12 – 15 MOS.	-	
VARICELLA	12-15 MOS.	4 WEEKS	Child <u>≥</u> 13 years-2 doses, minimum 4 weeks interval
HEP. A #1	<u>≥</u> 24 MOS.	6 MOS.	2 <sup>nd</sup> dose 6 to 12 mos.
HEP. A #2	<u>≥</u> 30 MOS.	-	after 1 <sup>st</sup> .
Td #1	<u>≥</u> 7 Yrs. of age	4 WEEKS	Booster every 10
Td #2		6 MOS.	years.
Td #3			
INFLUENZA #1	6 MOS.	4 WEEKS	Recomm. 6 mos. – 9 yrs.
INFLUENZA #2	-	-	first time flu shot
Pneumococcal Polysaccharide(PPV)	2 YEARS	5 YEARS	2 <sup>ND</sup> Dose only for child at highest risk of disease

+ This Hib dose is administered if you are administering a 4 dose Hib, or the child previously had one dose of the 4 dose Hib.

! Hep B & Hib administered separately, not as Comvax.

!! Infant must be ≥6 months of age before Hep.B is series complete (Hep B alone, not Comvax).